
Intellectual Property Network for Business

S.

IPN Home | Search | Order | Shopping Cart | Login | Help



WO9839293A2: 13-THIA PROSTAGLANDINS FOR USE IN GLAUCOMA THERAPY

View Images (33 pages) | View Cart

Premium Data 1: PDF (~3140 KB) | TIFF (~2480 KB) | Fax | More choices...



Issued/Filed Dates

pplication Number

Priority Number(s):

PC Class:

Designated:

Countries:

Abstract.

ZINKE, Paul, W., 4129 Willow Way Road, Fort Worth, TX 76133, United States of America

HELLBERG, Mark, R., 5211 Overridge Drive, Arlington, TX 76017, United States of America

ALCON LABORATORIES, INC., 6201 South Freeway, Fort Worth, TX 76134-2099, United States of America

Sept. 11, 1998 / March 6, 1998

WO1998US0004505

C07C 405/00;

March 7, 1997 US1997060040051

AU, BR, CA, CN, JP, KR, MX, US, European patent: AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE

13-thia prostaglandins are useful in the treatment of glaucoma and ocular hypertension. Also disclosed are ophthalmic, pharmaceutical compositions comprising said prostaglandins.

[Show "fr" Abstract]

Attorney, Agent, or Firm: Foreign References: COPELAND, Barry, L.;

none


(No patents reference this one)

Rowered by DB2 and Net Data

Nominate this invention for the Gallery...

Alternate Searches











SEARCH PATENT FULL TEXT WITH NATURAL LANGUAGE

¹ **Premium Data:** Using the premium data links may invoke a charge to your account. You can check the <u>pricelist here.</u>

Legal | IBM | FAQ | Feedback | Contact Us

Use of 13-thiaprostaglandin derivatives - for the treatment of gleucoma and ocular hypertension.

Drug Activity: Ophthalmological; Hypotensive

Mechanism of Action: Prostaglandin Compound Name: None Given

Use: For the treatment of glaucoma and ocular hypertension (claimed).

Dosage: 0.01-1000 (0.1-100) µg/eye topically.

Advantage: Reduced side effects, increased discrimination amongst receptors, improved therapeutic profile. Biological Data: (Ia) was tested for its intraocular pressure (IOP) lowering effect in cynomologus monkey eyes where ocular hypertension had been induced. Baseline IOP values were determined prior to treatment and 16 hours after the fourth dose. 3.0 mg of (Ia) caused an 18 +/- 3.0% reduction in IOP, compared to a 5.8 +/- 4.0% reduction achieved under identical conditions using PGF2α. Also presented is data showing that (Ia) causes less conjunctival hyperemia, conjunctival swelling and discharge than PGF2β.

<u>Chemistry</u>: The use of 13-thia prostaglandins of formula (I) is claimed for the treatment of glaucoma and ocular hypertension.

R1 = CO2R, ester, CONR4R5, CH2OR6 or CH2NR7R8; R = H or a cationic salt thereof; R4, R5 = H or alkyl. R6 = H, acyl or alkyl; R7, R8 = H, acyl, or alkyl provided that if either R7 or R8 = acyl, then the other = H or alkyl; n = 0 or 2; R2, R3 = H, alkyl or acyl; B = H, and OH in either configuration, H and F in either configuration, double bonded O, or OCH2CH2O; X = (CH2)q or (CH2)qO; q = 1-6; Y = 1-6C alkyl group, or a phenyl ring (optionally substituted); or X-Y = (CH2)pY1; p = 0-6; Y1 = further defined aromatic moiety; <math>a = single or double bond.

(I) is e.g. (5Z)-(9S, 11R, 15S)-9,11,15-trihydroxy-16-m-chlorophenoxy-13-thia-17,18,19,20-tetranor-5-prostenoic acid isopropyl ester (Ia) (Example II).

33 pages

Drawings 0/0

Authors: Zinke PW; Hellberg MR Publication Date: 11 September 1998

Language: English

Priority: 07 March 1997 US-040051

Location: Fort Worth, Tex., USA Document Number: WO9839293-A2 Filed: 06 March 1998 as U04505

Designated States: Regional: AT BE CH DE DK ES FI FR GB GR IE IT LU MC NL PT SE National: AU BR

CA CN JP KR MX US

WD-98-011852

PP - Cardiovascular

Page - 44